



Screening of Catalysts for Bio-oil Hydrodeoxygenation

Mortensen, Peter Mølgaard; Jensen, Anker Degn; Jensen, Peter Arendt

Publication date:
2012

[Link back to DTU Orbit](#)

Citation (APA):

Mortensen, P. M., Jensen, A. D., & Jensen, P. A. (2012). *Screening of Catalysts for Bio-oil Hydrodeoxygenation*. Abstract from 15th International Congress on Catalysis, Munich, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Screening of Catalysts for Bio-oil Hydrodeoxygenation

*Peter M. Mortensen, Anker D. Jensen, Peter A. Jensen, Department of Chemical and Biochemical Engineering, Technical University of Denmark, Lyngby, Denmark;
Jan-Dierk Grunwaldt, Institute of Chemical Technology and Polymer Science,
Karlsruhe Institute of Technology, Karlsruhe, Germany*

Introduction

A prospective route for production of bio-fuels is the conversion of biomass into bio-oil through pyrolysis followed by upgrading via hydrodeoxygenation (HDO) [1]. Through the pyrolysis, the bulk biomass is converted into a more energy-dense oil, called bio-oil. This is a viscous, polar, and acidic oil with a low heating value and high chemical instability, making it unsuitable as a fuel. These unfavourable characteristics are all associated with high contents of water (10-30 wt%) and oxygen containing organic compounds (30-40 wt% oxygen) in the oil [2]. In HDO the bio-oil is treated with hydrogen at a pressure of up to 200 bar and temperatures in the range 200-400 °C. This converts the oxygenates to a hydrocarbon product which will separate from the water and ultimately give a product equivalent to crude oil [1].

One of the major challenges in this concept is to find a catalyst for HDO which has a high activity for the deoxygenation reaction, but at the same time a sufficient lifetime [1]. A catalyst for HDO has to be bifunctional as both an activation of the oxy-compound and a donation of hydrogen is necessary for the reaction to take place [3]. Depending on the type of catalyst, these two steps can occur differently.

In the current work, three different types of catalysts for HDO have been tested: oxide catalysts, methanol synthesis catalysts, and reduced metal catalysts.

Experimental

All catalysts were prepared by the incipient wetness technique. Hereafter they were dried at 100 °C over night and calcined at 400°C for 4 h. Catalyst testing was performed in a 300ml batch reactor (Parr 4566). Analysis of the product was made on a Shimadzu GCMS/FID-QP2010UltraEi fitted with a Supelco Equity-5 column.

Results & Discussion

Figure 1 summarizes the results from the tested catalysts. A very low activity was found for the oxide and methanol synthesis type catalysts and neither of these achieved a conversion above 10%.




Figure 1 Yield and conversion over different types of catalysts. The experiments were conducted with 0.5 g of catalyst in 90 ml water and 10 g phenol. T=275 °C, P=100 bar, t=4 h.

Reduced metals are in Figure 1 represented by Ni, but were also tested in the combination with oxides. It was found that Ni based catalysts appeared to have a high activity relative to the other systems tested. Furthermore it was found that a system of Ni in combination with V_2O_5 had an even higher activity for HDO compared to Ni alone, as a much higher yield of cyclohexane was observed.

Conclusions

These results show that none of the here tested oxide type nor methanol synthesis type catalysts have a significant activity for HDO. Instead, Ni based catalysts appear very attractive and especially in the combination with an oxide, as V_2O_5 .

References

- [1] P.M. Mortensen, J.-D. Grunwaldt, P.A. Jensen, A.D. Jensen, A Review of catalytic upgrading of bio-oil to engine fuels, *Appl. Catal. A*, 407, 2011.
- [2] Q. Zhang, J. Chang, T. Wang, Y. Xu, Review of biomass pyrolysis oil properties and upgrading research, *Energy Convers. Manage.*, 48, 2007.
- [3] V.A. Yakovlev, et al., Development of new catalytic systems for upgraded bio-fuels production from bio-crude-oil and biodiesel, *Catal. Today*, 144, 2009.